

# Prevalence of GIT Nematodes and Associated Risk Factors of Exotic Chickens in Selected Farm of Poultry in and Around Ambo, Ethiopia

Abraham Belete Temesgen<sup>1\*</sup> , Zerihun Getie Wassie<sup>2</sup>  and Saleamlak Abebe<sup>3</sup>

<sup>1</sup>Department of Veterinary Pathobiology, College of Veterinary Medicine and Animal Sciences, University of Gondar, Ethiopia

<sup>2</sup>Department of Biomedical Science, College of Veterinary Medicine and Animal Sciences, University of Gondar, Ethiopia

<sup>3</sup>Department of School of Veterinary Medicine, College of Agriculture, Woldia University, Ethiopia

## \*Corresponding Author

Abraham Belete Temesgen, Department of Veterinary Pathobiology, College of Veterinary Medicine and Animal Sciences, University of Gondar, Ethiopia.

**Email:** abraham.belete@uog.edu.et

**Submitted:** 2024, Jun 27; **Accepted:** 2024, July 17; **Published:** 2024, Sep 03

**Citation:** Temesgen, A. B., Wassie, Z. G., Abebe, S. (2024). Prevalence of GIT Nematodes and Associated Risk Factors of Exotic Chickens in Selected Farm of Poultry in and Around Ambo, Ethiopia. *J Emerg Med OA*, 2(1), 01-06.

## Abstract

Poultry are raised worldwide in backyards and commercial systems with fewer social and religious taboos than other livestock. However, the chicken industry faces significant challenges from nematode parasites. A cross-sectional study with a random sampling technique was conducted from April to June 2019 to estimate gastrointestinal (GI) nematode parasites in chickens in selected farms in and around Ambo, Ethiopia. The fecal analysis results revealed that out of 70 samples collected, 60% were positive for gastrointestinal (GI) nematode eggs. Prevalence varied significantly by body condition, with the highest rates in chickens in poor condition (78.57%), followed by medium (54.54%) and good condition (40%). Location also played a significant role, with Ambo University Poultry Farm having the highest prevalence (83.87%), followed by Abebe Private Farm (55%) and Guder Campus Poultry Farm (26.31%). The main nematode species identified were *Ascaridia galli* (57.1%) and *Heterakis gallinarum* (2.9%). Infestation rates differed significantly by sex, age, location, and body condition, with males having higher rates of *Ascaridia galli* (61.53%) than females (56.14%), and *Heterakis gallinarum* exclusively affecting females (3.51%). Adults showed significantly higher rates of *Ascaridia galli* (85.71%) than young chickens (38.09%), with some infestation of *Heterakis gallinarum* (7.14%) observed in adults but absent in young chickens. This prevalence rate suggests limited awareness among chicken producers and insufficient control strategies in the study area. Hence, implementing targeted control strategies is advisable.

**Keywords:** Ambo, Chicken, Gastrointestinal Nematode, Prevalence

## 1. Introduction

Poultry are kept in backyards or commercial production systems in most areas of the world. Compared to a number of other livestock species, fewer social and religious taboos are related to the production, marketing, and consumption of poultry products. For these reasons, poultry products have become one of the most important protein sources for humans throughout the world [1]. In developing countries, poultry production offers an opportunity to feed the fast-growing human population and to provide income resources for poor farmers. Moreover, poultry in many parts of the modern world are considered as the chief source of not only cheaper protein of animal origin but also high-quality human food [2].

Among the important species of livestock kept in Ethiopia, poultry production systems are identified in the country. These include backyard poultry production systems, small-scale, and large-scale intensive production systems [3]. The population of poultry in Ethiopia is estimated to be 44.89 million, excluding the pastoral and agro-pastoral areas. With regard to breed, 96.46%, 0.57%, and 2.97% of the total poultry are reported to be indigenous, hybrid, and exotic, respectively [4]. Despite the presence of a large number of chickens in Ethiopia, the contribution to the national economy or the exploited benefit is very limited due to nutritional limitations and diseases [5]. Among parasitic diseases of poultry, nematode parasites are one of the major problems of the chicken industry

worldwide, characterized by ruffled feathers, loss of appetite, poor growth, and reduced egg production [6]. Moreover, nematodes (roundworms) are the most important group of helminth parasites of poultry. This is due to the large number of parasitic species that cause damage to the host, especially in severe infections. Most roundworms affect the gastrointestinal tract, with occasional parasites affecting the trachea or eye. Each species of roundworm tends to infect a specific area of the gastrointestinal tract. Different species of the same genus may infect several different areas of the tract. In general, the different species of roundworms have very similar life cycles [7]. Of the helminth parasites of poultry birds, nematodes constitute the most important group of helminth parasites of poultry, both in the number of species and the extent of damage they cause; the main genera include *Ascaridia* and *Heterakis* [8].

Generally, nematode infections in poultry are widely distributed in different parts of the world, and numerous research efforts have been undertaken to prevent poultry mortality from parasitic diseases. The prevalence of two nematode genera, *Ascaridia* and *Heterakis*, has been extensively studied [9]. Among diseases, internal parasites are known to reduce the productivity of poultry kept under various management systems. Infection by parasites occurs after ingestion of nematode eggs or intermediate hosts such as cockroaches, grasshoppers, ants, and earthworms. Nematode infection results in a reduction in food intake, injury to the intestinal

wall, and hemorrhage, leading to poor absorption of nutrients and weight loss [10]. Gastrointestinal parasite infestation is a common problem in poultry, especially when nematode infections occur in high proportions in animals reared in intensive management systems. In Ethiopia, the poultry industry is developing for both local and exotic chickens, but only a few surveys have been carried out to determine the burden of nematode parasites in chickens in this country [11,12]. Therefore, the objectives of this study were to estimate the prevalence of major gastrointestinal nematode parasites in poultry and to assess the risk factors associated with the incidence of the parasites in the study area.

## 2. Materials and Methods

### 2.1 Study Area

The study was conducted on selected farms in and around Ambo, Ethiopia (Figure1), from April to June 2019. Ambo is the administrative center of the zone and is located at a latitude and longitude of 8°59'N 37°51'E, with an elevation of 2101 meters above sea level (asl) and 114 km west of Addis Ababa. The Ambo Woreda has 34 administrative kebeles. The agro-ecology of the study area is 23% highland, 60% midland, and 17% lowland. It has an annual rainfall and temperature ranging from 800-1000 mm and 20-29 °C, respectively. The livestock population of the district includes 145,371 cattle, 50,152 sheep, 27,026 goats, 105,794 chickens, 9,088 horses, 2,914 donkeys, and 256 mules [13].




Figure 1: Map of the Study Area

---

## 2.2 Study Population

The study populations included all exotic breeds' chickens of both sexes and all age groups that were kept under intensive management systems from the selected intensive farms. The birds were categorized into two age groups: Young (under 12 months) and adult (above 12 months of age). The age of the birds was determined from records found on the farm and subjectively based on the size of the crown, length of the spur, and so on [14].

## 2.3 Study Design and Sample Size Determination

A cross-sectional study design was used for this study. Sex, different age groups, body condition, breed, and management systems were recorded as test variables during the data collection of target chickens. A random sampling technique was used to recruit animals for the study. The required sample size for this study was estimated according to the formula of, with an expected prevalence of 68.5% from a previous study conducted by in a comparable agro-ecological area, and a desired absolute precision ( $d$ ) of 0.05 at a 95% confidence level [12,15].

$$\frac{1.96^2 \times P_{exp} \times (1 - P_{exp})}{d^2}$$

Where  $n$  represents the required sample size,  $P_{exp}$  denotes the expected prevalence, and  $d$  stands for the desired absolute precision. Using this formula, the sample size was calculated to be 332. However, only 70 chickens were selected purposefully due to a shortage of time.

## 3. Data Collection

After examining the selected chickens for general body condition and clinical signs indicative of gastrointestinal nematodes, samples were taken and entered into the Ambo University Department of Veterinary Laboratory Technology laboratory for parasitological examinations. Fecal samples were collected using hand gloves directly from the vent or top surface of freshly voided feces and placed in airtight screw-cap universal bottles. The samples were transported to the Parasitology laboratory of Ambo University, College of Veterinary Laboratory Technology, and stored at 4°C until examination. During sample collection, information about risk factors such as body condition, sex, age, and all other management systems were recorded for each collected sample. Samples were processed using the floatation technique as described by [16].

## 4. Data Management and Analysis

The information obtained from laboratory tests and observations was entered into the spreadsheet of Microsoft Excel worksheet. Descriptive statistics were analyzed using SPSS version 20 software, and the Chi-square ( $\chi^2$ ) test was used to analyze the sample data. The Chi-square test was used to assess whether there is a statistically significant difference in gastrointestinal nematode infection between sexes and ages. A statistically significant association between variables was considered to exist if the calculated p-value is less than 0.05 with a 95% confidence level.

## 5. Results

The fecal analysis results revealed that out of 70 samples collected, 42 (60%) were positive for gastrointestinal (GI) nematode eggs. The prevalence of nematode infection varied significantly with body condition: chickens in poor body condition had the highest prevalence at 78.57%, followed by those in medium condition at 54.54%, and those in good condition at 40%. This difference was statistically significant. Location also played a significant role in the prevalence of GI nematodes. The highest prevalence was observed at Ambo University Poultry Farm (83.87%), compared to Abebe Private Farm (55%) and Guder Campus Poultry Farm (26.31%), with the differences being statistically significant. Among the risk factors evaluated, age, body condition, and location showed statistically significant differences in prevalence, whereas sex did not exhibit a significant difference ( $p > 0.05$ ) (Table 1 and Figure 2).


The overall prevalence of two nematode species identified from fecal examinations in chickens were *Ascaridia galli* (57.1%) and *Heterakis gallinarum* (2.9%). The infestation rates of these nematodes vary significantly based on sex, age, location, and body condition. Males have a slightly higher infestation rate of *Ascaridia galli* (61.53%) compared to females (56.14%), while *Heterakis gallinarum* is only observed in females (3.51%). Age-wise, adults exhibit a significantly higher infestation rate of *Ascaridia galli* (85.71%) compared to young chickens (38.09%) and also show some infestation of *Heterakis gallinarum* (7.14%), which is absent in young chickens. Location-wise, chickens from Ambo University have the highest infestation rate of *Ascaridia galli* (77.41%) and a noticeable infestation of *Heterakis gallinarum* (6.45%), whereas those from Guder Campus have the lowest rates for both parasites (26.31% and 0%, respectively). Chickens in poor body condition are the most affected, with the highest infestation rates for *Ascaridia galli* (71.42%) and *Heterakis gallinarum* (7.14%), while those in good condition have the lowest rates for both nematodes (Table 2 and Figure 3).

Factors	Category	Examined	Positive	$\chi^2$	P-value
Sex	Female	57	34 (59.64%)	0.016	0.900
	Male	13	8 (61.53%)		
Age	Young	42	16 (38.09%)	20.992	0.000
	Adult	28	26 (92.85%)		
	Abebe private	20	11 (55%)		
Place (Farm)	Ambo university	31	26 (83.87%)	16.55	0.000
	Guder campus	19	5 (26.31%)		
Body condition	Poor	28	22 (78.57%)	7.630	0.022
	Medium	22	12 (54.54%)		
	Good	20	8 (40%)		
Total		70	42(60.0%)		


**Table 1: The Prevalence Gastro-Intestinal Nematode Infection by Risk Factors**

Factors	Category	Examined	Nematode spp		$\chi^2$	P-value
			<i>Ascaridia galli</i>	<i>Heterakis Gallinarum</i>		
Sex	Female	57	32 (56.14%)	2 (3.51%)	0.520	0.771
	Male	13	8 (61.53%)	0 (0%)		
Age	Young	42	16 (38.09%)	0 (0%)	22.262	0.000
	Adult	28	24 (85.71%)	2 (7.14)		
	Abebe private	20	11 (55%)	0 (0%)		
Place (Farm)	Ambo university	31	24 (77.41%)	2 (6.45%)	17.853	0.001
	Guder campus	19	5 (26.31%)	0(0%)		
Body condition	Poor	28	20 (71.42%)	2 (7.14%)	9.632	0.047
	Medium	22	12 (54.54%)	0 (0%)		
	Good	20	8 (40%)	0 (0%)		
Total		70	40(57.1%)	2(2.9%)		

**Table 2: The Prevalence of Nematode Species Infestation by Risk Factors**



**Figure 2: The Prevalence of Nematode Species Infestation by Risk Factors**



**Figure 3: The Prevalence of Nematode Species Infestation by Risk Factors**

## 6. Discussion

In this cross-sectional study, the overall prevalence of infection with gastrointestinal (GI) nematode eggs was 60.0%. This finding was higher than previous reports by, who reported a prevalence of 59.64% and 53% in Ethiopia and Nigeria, respectively [8]. Nonetheless, this finding was lower than the report of, which reported 61.9%, and who reported 64.7% in Nigeria and in the Oromia region of Ethiopia, respectively [17,18]. Additionally, the current study was slightly lower than the result recently reported by of 68.5% from the same area in Ambo, West Shoa Zone, Oromia Regional State, Ethiopia [12]. This discrepancy could be related to differences in the management system, study method, sample size, and control practices in the area. In the present study, females seemed to have a higher prevalence (59.64%) than males (61.53%), which could be related to the higher susceptibility of female animals. However, there was no significant difference ( $\chi^2 = 0.016$  and  $P > 0.05$ ) in the prevalence of gastrointestinal nematode parasites between sexes. This finding aligns with, who reported that female chickens were more infected with GI nematode parasites than males [8,19]. Female chickens are known to be more voracious in their feeding habits, especially during egg production, compared to males. However, this study contrasts with another report from Haromaya by, found higher GI nematode infection in males (52.1%) than females (39.9%) [20]. This difference may be due to sample size and nutritional deficiency. Another report by indicated no usual natural affinity of GI nematode species to either sex of the host chickens [21].

Among the age groups, adult chickens had a higher prevalence (92.85%) than young age groups (38.09%), with a statistically significant difference ( $p < 0.05$ ). This could be due to adult chickens being exposed to infective larval stages for a longer time, contributing to a higher prevalence in older age groups. This finding is consistent with but contrary to, who reported a higher level of GI nematode prevalence in young chickens [16,22]. This

could be due to young chickens having a lower level of immunity compared to adults. Similarly, observed a higher prevalence of GI nematodes in young chickens than in adults in Kenya [23]. The most prevalent nematode species encountered in the present study was *Ascaridia galli* (57.1%). The prevalence of *Ascaridia galli* was higher than the previously reported works in central Ethiopia by (55.26%) and from Haromaya (38.0%) [20,24]. This might be due to differences in management systems, deworming practices, and/or agro-ecological conditions of the study area. The high occurrence of the parasites in the study area may be related to the wet season when the survey was conducted. The prevalence of *H. gallinarum* (2.9%) in this study was lower than in another study in Ethiopia (4.3%) and higher than another study in Kenya (1.43%) [25,26]. However, it is lower than the 51.6% reported by in Ethiopia, possibly due to agro-ecological variation [27].

## 7. Conclusion

The current study indicated that the overall prevalence of gastrointestinal nematode (GIT) infections in chickens in the study area was 60.0%. GIT nematode infections are found to be an important problem in the study area. The results of this study showed that *Ascaridia galli* was the most dominant species. The compared risk factors in the current study, such as body condition, place, and age, showed significant variation with nematode infection. This prevalence rate suggests limited awareness among chicken producers and insufficient control strategies in the study area. Therefore, implementing targeted control strategies is advisable.

## Data Sharing Statement

All data generated or analyzed during this study are available upon request from the corresponding author.

## Acknowledgement

The author expresses heartfelt gratitude to all co-authors of this

manuscript for their invaluable contributions and support.

### Funding

None

### Disclosure

The author declares no conflicts of interest in this work.

### References

1. Alders, R. G. (2004). *Poultry for profit and pleasure* (No. 3). Food & Agriculture Org..
2. Central Statistical Agency (CSA) (2009). Volume II: Report on livestock and livestock characteristics (private peasant holdings) (Statistical Bulletin 468). Addis Ababa: Federal Democratic Republic of Ethiopia.
3. Yami A, Tadella D. (1997). The status of poultry research and development in Ethiopia. *D.Z.A.R.C. Res Bull.* 4:40-46.
4. Central Statistical Agency (CSA). (2012). Volume II: Report on livestock and livestock characteristics (private peasant holdings) (Statistical Bulletin 570). Addis Ababa: Federal Democratic Republic of Ethiopia.
5. Smith A.J. (1990). *Poultry: The tropical agriculturalist*. Hong Kong: Macmillan Publisher.
6. Aiello S, Mays A. (1998). *The Merck veterinary manual*. 8th ed. Whitehouse Station, NJ: Merck and Co. Inc.
7. Belete, A., Addis, M., & Ayele, M. (2016). Review on major gastrointestinal parasites that affect chickens. *Journal of Biology, Agriculture and Healthcare*, 6(11), 11-21.
8. Matur, B. M., Dawam, N. N., & Malann, Y. D. (2010). Gastrointestinal helminth parasites of local and exotic chickens slaughtered in Gwagwalada, Abuja (FCT), Nigeria. *New York Science Journal*, 3(5), 96-99.
9. Mulugeta, A., Chanie, M., & Bogale, B. (2013). Major constraints of village poultry production in Demba Gofa District of Southern Region, Ethiopia. *British Journal of Poultry Sciences*, 2(1), 01-06.
10. Soulsby E.J.L. (1986). *Helminthes, arthropods and protozoa of domesticated animals*. 7th ed. London: Bailliere Tindall.
11. Oljira, D., Melaku, A., & Bogale, B. (2012). Prevalence and risk factors of coccidiosis in poultry farms in and around Ambo Town, Western Ethiopia. *American-Eurasian Journal of Scientific Research*, 7(4), 146-149.
12. Shiferaw, S., Tamiru, F., Gizaw, A., Atalel, D., Terfa, W., Dandecha, M., & Mekibib, A. (2016). Study on prevalence of helminthes of local backyard and exotic chickens in and around Ambowest Shoa Zone, Oromia Regional State, Ethiopia. *J Veter Sci Med*, 4(2), 4.
13. Ambo Town Ministry of Agricultural Office (ATMA). (2010). Annual report. Ambo, Ethiopia.
14. Damerow, G. (1995). *Story's guide to raising chickens*. USA: Story Books.
15. Thrusfield, M. (1995). *Veterinary epidemiology*. Oxford: Blackwell Science.
16. Permin, A., & Hansen, J. W. (1998). Diagnostic methods: Epidemiology, diagnosis and control of poultry parasites. *FAO Animal Health Manual*, No 4. Rome: Food and Agriculture Organization of the United Nations
17. Luka, S. A., & Ndams, I. S. (2007). Short communication report: Gastrointestinal parasites of domestic chicken *Gallus-gallus domesticus linnaeus 1758* in Samaru, Zaria Nigeria. *Science World Journal*, 2(1).
18. Tolossa, Y., Shafi, Z., & Basu, A. (2009). Ectoparasites and gastrointestinal helminths of chickens of three agro-climatic zones in Oromia Region, Ethiopia. *Animal Biology*, 59(3), 289-297.
19. Sonaiya, E. B. (1990, October). The context and prospects for development of smallholder rural poultry production in Africa. In *CTA-Seminar proceedings on smallholder rural poultry production* (pp. 9-13).
20. Tesfaheywet, Z., Amare, E., & Hailu, Z. (2012). Helminthosis of chickens in selected small scale commercial poultry farms in and around Haramaya Woreda, Southeastern Ethiopia. *Journal of Veterinary Advances*, 2(9), 462-468.
21. Hassouni, T., & Belghyti, D. (2006). Distribution of gastrointestinal helminths in chicken farms in the Gharb region—Morocco. *Parasitology research*, 99, 181-183.
22. Mekuria, S., & Bayessa, M. (2017). Gastrointestinal helminths and their predisposing factors in different poultry management systems; Haromaya, Ethiopia. *Ethiopian Veterinary Journal*, 21(1), 40-43.
23. Mungube, E. O., Bauni, S. M., Tenhagen, B. A., Wamae, L. W., Nzioka, S. M., Muhammed, L., & Nginyi, J. M. (2008). Prevalence of parasites of the local scavenging chickens in a selected semi-arid zone of Eastern Kenya. *Tropical Animal Health and Production*, 40, 101-109.
24. Ashenafi H, Eshetu, Y. (2004). Study on gastrointestinal helminthes of local chickens in central Ethiopia. *J Vet Med*. 155:504-507.
25. Eshetu, Y., Mulualem, E., Ibrahim, H., Berhanu, A., Aberra, k. (1999). Study of gastrointestinal helminthes of scavenging chickens in four rural districts of Amhara. *Ethiopian Agricultural Research Organization (EARO), National Poultry Research Program: Strategy Document*. Addis Ababa, Ethiopia.
26. Kaingu, F. B., Kibor, A. C., Shivairo, R., Kutima, H., Okeno, T. O., Waihenya, R., & Kahi, A. K. (2010). Prevalence of gastro-intestinal helminthes and coccidia in indigenous chicken from different agro-climatic zones in Kenya. *African journal of Agricultural research*, 5(6), 458-462.
27. Berhanu, M., Haileyesus, D., Desie, S. (2014). Gastrointestinal helminthes of scavenging chickens in outskirts of Hawassa, Southern Ethiopia. *Global Veterinaria*, Vol. 12, 557-561.

**Copyright:** ©2024 Abraham Belete Temesgen, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.